

ON SOME QUESTIONS OF EYMARD AND BEKKA CONCERNING AMENABILITY OF HOMOGENEOUS SPACES AND INDUCED REPRESENTATIONS

VLADIMIR PESTOV

ABSTRACT. Let $F \subseteq H \subseteq G$ be closed subgroups of a locally compact group. In response to a 1972 question by Eymard, we construct an example where the homogeneous factor space G/F is amenable in the sense of Eymard–Greenleaf, while H/F is not. (In our example, G is discrete.) As a corollary which answers a 1990 question by Bekka, the induced representation $\text{ind}_H^G(\rho)$ can be amenable in the sense of Bekka even if ρ is not amenable. The second example, answering another question by Bekka, shows that $\text{ind}_H^G(\rho)$ need not be amenable even if both the representation ρ and the coset space G/H are amenable.

Résumé. Soient $F \subseteq H \subseteq G$ deux sous-groupes fermés d'un groupe localement compact G . En réponse à une question posée en 1972 par Eymard, nous construisons un groupe discret G tel que l'espace homogène G/F est moyennable au sens d'Eymard et Greenleaf, bien que H/F n'est pas moyennable. On obtienne un corollaire qui répond à un problème posé par Bekka en 1990: une représentation induite $\text{ind}_H^G(\rho)$ peut être moyennable au sens de Bekka même si ρ n'est pas moyennable. Le deuxième exemple, qui répond à une autre question de Bekka, montre que $\text{ind}_H^G(\rho)$ n'est pas nécessairement moyennable même si la représentation ρ et l'espace homogène G/H sont l'un et l'autre moyennables.

1. INTRODUCTION

Let H be a closed subgroup of a locally compact group G . The homogeneous factor space G/H is *amenable in the sense of Eymard and Greenleaf* [5, 6], if $L^\infty(G/H)$ supports a G -invariant mean. If $H = \{e\}$, one obtains the classical concept of an amenable locally compact group.

A unitary representation ρ of a group G in a Hilbert space $\mathcal{H}$ is *amenable in the sense of Bekka* [1] if there exists a state, ϕ , on the algebra $\mathcal{B}(\mathcal{H})$ of bounded operators that is $\text{Ad } G$ -invariant: $\phi(\pi(g)T\pi(g)^{-1}) = \phi(T)$ for every $T \in \mathcal{B}(\mathcal{H})$ and every $g \in G$. For instance, the homogeneous space G/H is Eymard–Greenleaf amenable if and only if the quasi-regular representation $\lambda_{G/H}$ of G in $L^2(G/H)$ is amenable.

Let F and H be closed subgroups of a locally compact group G , such that $F \subseteq H \subseteq G$. In 1972 Eymard had asked ([5], p. 55):

2000 Mathematical Subject Classification. 43A07, 43A65, 43A85, 22D30.

Partially supported by the Marsden Fund of the Royal Society of New Zealand and by a University of Ottawa start-up grant.

Q 1. *Suppose the space G/F is amenable. Is then H/F amenable?*

This is of course a classical result in the case $F = \{e\}$.

Let π be a strongly continuous unitary representation of H . In 1990 Bekka has shown [1] that, if the unitarily induced representation $\text{ind}_H^G(\pi)$ is amenable, then G/H is amenable. He asked the following:

Q 2. *Does amenability of $\text{ind}_H^G(\pi)$ imply that of π ?*

(This is a more general version of Eymard's question Q 1.)

Q 3. *Suppose both G/H and π are amenable. Is then $\text{ind}_H^G(\pi)$ amenable?*

Question 3 was partially answered in the positive by Bekka himself [1] under extra assumptions on H . Further discussion can be found in [7].

We show that in general the answer to all three questions above is negative.

Remark. In a recent independent work Nicolas Monod and Sorin Popa have also solved Eymard's problem (and thus Bekka's Q 2 as well). (See their note [8] in the same issue.)

2. REMINDERS AND SIMPLE FACTS

2.1. A unitary representation π of a locally compact group G in a Hilbert space $\mathcal{H}$ is said to *almost have invariant vectors* if for every compact $K \subseteq G$ and every $\varepsilon > 0$ there is a $\xi \in \mathcal{H}$ satisfying $\|\xi\| = 1$ and $\|\pi_g(\xi) - \xi\| < \varepsilon$ for all $g \in K$.

2.2. *If a unitary representation π of a group G (viewed as discrete) almost has invariant vectors, then π is amenable.*

This follows from Corollary 5.3 of [1], because π weakly contains the trivial one-dimensional representation, $1_G < \pi$. Alternatively, choose for every finite $K \subseteq G$ and every $\varepsilon > 0$ an almost invariant vector $\xi_{K,\varepsilon}$ as above, and set $\phi_{K,\varepsilon}(T) := \langle T\xi_{K,\varepsilon}, \xi_{K,\varepsilon} \rangle$. Every weak* cluster point, ϕ , of the net of states $(\phi_{K,\varepsilon})$ is a G -invariant state.

Note that the converse of 2.2 is not true, as for example every unitary representation of an amenable group is amenable by Theorem 2.2 of [1].

2.3. *Let H be a closed subgroup of a locally compact group G . The following statements are equivalent.*

- (i) *The homogeneous space G/H is amenable.*
- (ii) *The left quasi-regular representation $\lambda_{G/H}$ of G in $L^2(G/H)$, formed with respect to any (equivalently: every) quasi-invariant measure ν on G/H , almost has invariant vectors.*
- (iii) *The left quasi-regular representation $\lambda_{G/H}$ is amenable.*
- (iv) *If G acts continuously by affine transformations on a convex compact set C in such a way that C contains an H -fixed point, then C contains a G -fixed point.*

The equivalences between (i), (ii), and (iv) are due to Eymard [5], while (iii) was added by Bekka (Th. 2.3.(i) of [1]). The condition (ii) is an analogue of Reiter's condition (P_2) for homogeneous spaces. (Interestingly, a natural analogue of Følner's condition for homogeneous spaces fails [6].)

2.4. *Let the homogeneous space G/H be amenable, and let π be a strongly continuous unitary representation of G . If the restriction of π to H is an amenable representation, then π is amenable as well.*

To prove this statement, denote, following Bekka (Section 3 of [1]), by $X(\mathcal{H})$ the C^* -subalgebra of $\mathcal{B}(\mathcal{H})$ formed by all operators T with the property that the orbit map of the adjoint action,

$$G \ni g \mapsto \pi(g)T\pi(g)^{-1} \in \mathcal{B}(\mathcal{H}),$$

is norm-continuous. Then G acts upon $X(\mathcal{H})$ in a strongly continuous way by isometries. It follows that the dual adjoint action of G on the state space of $X(\mathcal{H})$, equipped with the weak* topology, is continuous. Because of the assumed amenability of $\pi|_H$, there is an H -invariant state, ψ , on $\mathcal{B}(\mathcal{H})$ and consequently on $X(\mathcal{H})$. Eymard's conditional fixed point property 2.3.(iv) allows us to conclude that there is a G -invariant state, ϕ , on $X(\mathcal{H})$. This in turn implies amenability of π by Theorem 3.5 of [1].

2.5. Let H be a closed subgroup of a locally compact group G , and let π be a strongly continuous unitary representation of H in a Hilbert space. By $\text{ind}_H^G(\pi)$ we will denote, as usual, the unitarily induced representation of G .

2.6. The following is Mackey's generalization of the Frobenius Reciprocity Theorem (Theorem 5.3.3.5 of [9]).

Let H be a closed subgroup of a locally compact group G , and let π and ρ be finite-dimensional irreducible unitary representations of H and G , respectively. If G/H carries a finite invariant measure, then $\text{ind}_H^G(\pi)$ contains ρ as a discrete direct summand exactly as many times as $\rho|_H$ contains π as a discrete direct summand.

2.7. Let G be a locally compact group, and let H, F be closed subgroups of G such that $F \subseteq H$. Set $\pi = \text{ind}_F^H(1_F)$, where 1_F stands for the trivial one-dimensional representation of F . Then $\pi = \lambda_{H/F}$ is unitarily equivalent to the quasi-regular representation of H in $L^2(H/F)$ ([9], Corollary 5.1.3.6). By the theorem on induction in stages (*ibid.*, Proposition 5.1.3.5), the induced representation $\text{ind}_H^G(\lambda_{H/F})$ is unitarily equivalent to $\text{ind}_F^G(1_F)$, which is just the representation $\lambda_{G/F}$, the quasi-regular representation of G in $L^2(G/F)$.

Now assume that the homogeneous factor space G/F is amenable. By Bekka's result mentioned above (2.3.(iii)), this amounts to the amenability of $\lambda_{G/F}$.

This argument shows that Eymard's Q 1 is a particular case of Bekka's Q 2.

3. FIRST EXAMPLE

3.1. Let $H = F_\infty$, where F_∞ denotes the free non-abelian group on infinitely many free generators x_i , $i \in \mathbb{Z}$. For every $n \in \mathbb{Z}$, let Γ_n denote the normal subgroup of F_∞ generated by x_i , $i \leq n$. We will set $F = \Gamma_0$.

As the factor space $H/F = F_\infty/\Gamma_0$ is a free group, it is not amenable.

3.2. Let the group $\mathbb{Z}$ act on F_∞ by group automorphisms via shifting the generators:

$$\tau_n(x_m) := x_{m+n},$$

where $m, n \in \mathbb{Z}$ and τ denotes the action of $\mathbb{Z}$ on F_∞ .

Denote by $G = \mathbb{Z} \ltimes_\tau F_\infty$ the semi-direct product formed with respect to the action τ . That is, G is the Cartesian product $\mathbb{Z} \times F_\infty$ equipped with the group operation $(m, x)(n, y) = (m + n, x\tau_n y)$, the neutral element $(0, e)$ and the inverse $(n, x)^{-1} = (-n, \tau_{-n}x^{-1})$.

Let us show that the homogeneous space G/F is Eymard–Greenleaf amenable.

3.3. For every $n \in \mathbb{Z}$,

$$\begin{aligned} (n, e)F &= \{(n, e)(0, y) : y \in F\} \\ &= \{(n, \tau_n y) : y \in \Gamma_0\} \\ &= \{n\} \times \Gamma_n. \end{aligned}$$

Let $S \subset F_\infty$ be an arbitrary finite subset. For $n \in \mathbb{Z}$ sufficiently large, $S \subset \Gamma_n$.

Now for each $(0, s) \in S$,

$$\begin{aligned} (0, s)(n, e)F &= \{(0, s)(n, z) : z \in \Gamma_n\} \\ &= \{n\} \times (s\Gamma_n) \\ &= \{n\} \times \Gamma_n \\ &= (n, e)F; \end{aligned}$$

that is, the left F -coset $(n, e)F \in G/F$ is S -invariant.

Consequently, the unit vector $\delta_{(n, e)F} \in \ell^2(G/F)$ is S -invariant. We have proved that the H -module $\ell^2(G/F)$ almost has invariant vectors, and therefore the restriction to H of the left regular representation $\lambda_{G/F}$ is amenable. Since G/H is an amenable homogeneous space (H is normal in G and the factor-group G/H is isomorphic to $\mathbb{Z}$), we conclude by 2.4 that the left regular representation of G in $\ell^2(G/F)$ is amenable, that is, that G/F is an amenable homogeneous space.

3.4. The constructed triple of groups $F \subset H \subset G$ provides a negative answer to the question of Eymard (Q 1).

In view of the remarks in 2.7, it provides a negative answer to Bekka's question (Q 2) as well. Take as $\pi = \text{ind}_F^H(1_F) = \lambda_{H/F}$ the left quasi-regular representation of H in $\ell^2(H/F)$. While π is not amenable, the induced representation $\text{ind}_H^G(\pi) = \lambda_{G/F}$ is amenable.

3.5. The following way of viewing our example may be instructive.

If considered as a unitary F_∞ -module, $\ell^2(G/F)$ decomposes, up to unitary equivalence, into the orthogonal sum

$$\ell^2(G/F) \cong \bigoplus_{n \in \mathbb{Z}} \ell^2(F_\infty/\Gamma_n).$$

None of the unitary F_∞ -modules $\ell^2(F_\infty/\Gamma_n)$ is amenable, yet their orthogonal sum is amenable, because the family $(\ell^2(F_\infty/\Gamma_n))_{n \in \mathbb{Z}}$ “asymptotically has invariant vectors”: every finite $S \subset F_\infty$ acts trivially on the Hilbert space $\ell^2(F_\infty/\Gamma_n)$, provided n is large enough so that $S \subset \Gamma_n$. (Indeed, Γ_n are normal in F_∞ .)

4. SECOND EXAMPLE

4.1. Let $G = \mathrm{SL}(n, \mathbb{R})$ and $H = \mathrm{SL}(n, \mathbb{Z})$, $n \geq 3$. Since H is a (non-uniform) lattice in G (cf. Exercise 7, §2, Chapitre VII of [3]), the homogeneous space G/H is amenable.

4.2. The group $\mathrm{SL}(n, \mathbb{Z})$ is maximally almost periodic (for instance, homomorphisms to the groups $\mathrm{SL}(n, \mathbb{Z}_p)$, where p is a prime number, separate points in $\mathrm{SL}(n, \mathbb{Z})$). Let π be a non-trivial (of dimension > 1) irreducible unitary finite-dimensional representation of H . Being finite-dimensional, π is an amenable representation; cf. [1], Theorem 1.3.(i).

4.3. Applying Mackey’s Reciprocity Theorem (quoted above, 2.6) with ρ equal to 1_G , a trivial one-dimensional representation of G , we conclude that the unitarily induced representation $\mathrm{ind}_H^G(\pi)$ has no non-zero invariant vectors.

At the same time, the Lie group $G = \mathrm{SL}(n, \mathbb{R})$, $n \geq 3$, has property (T) (see *e.g.* [4], Chapitre 2.a, Théorème 4), and the only amenable representations of non-compact simple Lie groups with Kazhdan’s property (T), such as G , are those having a non-zero invariant vector. ([1], Remark 5.10; cf. also the principal result of [2].)

4.4. We conclude: $\mathrm{ind}_H^G(\pi)$ is non-amenable, even if the unitary representation π and the homogeneous space G/H are both amenable.

This answers in the negative another question posed by Bekka (Q 3).

REFERENCES

- [1] M.E.B. Bekka, *Amenable unitary representations of locally compact groups*, Invent. Math. **100** (1990), 383–401.
- [2] M.E.B. Bekka and A. Valette, *Kazhdan’s property (T) and amenable representations*, Math. Zeit. **212** (1993), 293–299.
- [3] N. Bourbaki, *Intégration*, Chapitres 7 et 8, Hermann, Paris, 1963.
- [4] P. de la Harpe and A. Valette, *La propriété (T) de Kazhdan pour les groupes localement compacts*, Astérisque **175** (1989), 158 pp.
- [5] P. Eymard, *Moyennes invariantes et représentations unitaires*, Lecture Notes Math. **300**, Springer-Verlag, Berlin-New York, 1972, 113 pp.
- [6] F.P. Greenleaf, *Amenable actions of locally compact groups*, J. Funct. Anal. **4** (1969), 295–315.

- [7] P. Jolissaint, *Invariant states and a conditional fixed point property for affine actions*, Math. Ann. **304** (1996), 561–579.
- [8] N. Monod and S. Popa, *On co-amenability for groups and von Neumann algebras*, this issue.
- [9] G. Warner, *Harmonic Analysis on Semi-Simple Lie Groups I*, Grundlehren der math. Wissenschaften Bd. **188**, Springer-Verlag, Berlin-Heidelberg-New York, 1972.

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, 585 KING EDWARD AVE., OTTAWA, ONTARIO, CANADA K1N 6N5.

E-mail address: vpest283@science.uottawa.ca